Food and Drug Administration, HHS

intended as source material for further manufacturing use. The definition excludes single donor plasma products intended for intravenous use.

[41 FR 10768, Mar. 12, 1976, as amended at 50 FR 4140, Jan. 29, 1985]

§ 640.61 Informed consent.

The written consent of a prospective donor shall be obtained after a qualified licensed physician has explained the hazards of the procedure to the prospective donor. The explanation shall include the risks of a hemolytic transfusion reaction if he is given the cells of another donor, and the hazards involved if he is hyperimmunized. The explanation shall consist of such disclosure and be made in such a manner that intelligent and informed consent be given and that a clear opportunity to refuse is presented.

EFFECTIVE DATE NOTE: At 80 FR 29905, May 22, 2015, §640.61 was removed, effective May 23, 2016

§ 640.62 Medical supervision.

A qualified licensed physician shall be on the premises when donor suitability is being determined, immunizations are being made, whole blood is being collected, and red blood cells are being returned to the donor.

[66 FR 1837, Jan. 10, 2001]

EFFECTIVE DATE NOTE: At 80 FR 29905, May 22, 2015, $\S640.62$ was removed, effective May 23, 2016.

§ 640.63 Suitability of donor.

- (a) Method of determining. The suitability of a donor for Source Plasma shall be determined by a qualified licensed physician or by persons under his supervision and trained in determining donor suitability. Such determination shall be made on the day of collection from the donor by means of a medical history, tests, and such physical examination as appears necessary to the qualified licensed physician.
- (b) Initial medical examinations. (1) Each donor shall be examined by a qualified licensed physician on the day of the first donation or no more than 1 week before the first donation and at subsequent intervals of no longer than 1 year.

- (2)(i) A donor who is to be immunized for the production of high-titer plasma shall be examined by a qualified licensed physician. The medical examination shall be performed within no more than 1 week before the first immunization injection. The medical examination for plasmapheresis need not be repeated, if the first donation occurs within 3 weeks after the first injection.
- (ii) A donor who is an active participant in a plasmapheresis program, and has been examined in accordance with paragraph (b)(1) of this section, need not be reexamined before immunization for the production of high-titer plasma.
- (3) Each donor shall be certified to be in good health by the examining physician. The certification of good health shall be on a form supplied by the licensed establishment and shall indicate that the certification applies to the suitability of the individual to be a plasmapheresis donor and, when applicable, an immunized donor.
- (c) Qualification of donor. Donors shall be in good health on the day of donation, as indicated in part by:
 - (1) Normal temperature;
- (2) Demonstration that systolic and diastolic blood pressures are within normal limits, unless the examining physician is satisfied that an individual with blood pressures outside these limits is an otherwise qualified donor under the provisions of this section:
- (3) A blood hemoglobin level of no less than 12.5 grams of hemoglobin per 100 milliliters of blood or a hematocrit level of 38 percent;
 - (4) A normal pulse rate;
- (5) A total serum or total plasma protein of no less than 6.0 grams per 100 milliliters of blood;
- (6) Weight, which shall be at least 110 pounds:
- (7) Freedom from acute respiratory diseases;
- (8) Freedom from any infectious skin disease at the site of phlebotomy and from any such disease generalized to such an extent as to create a risk of contamination of the plasma;
- (9) Freedom from any disease, other than malaria, transmissible by blood